

Toward the analogue of a thermally generated electromagnetic field

Bob Osano^{1,2,*} and Patrick W. M. Adams^{1,†}

¹*Astrophysics, Cosmology and Gravity Centre, Department of Mathematics and Applied Mathematics,
University of Cape Town, Rondebosch 7701, Cape Town, South Africa*

²*Academic Development Programme, Science, Centre for Higher Education Development,
University of Cape Town, Rondebosch 7701, Cape Town, South Africa*

(Dated: June 13, 2016)

Magnetic and vorticity fields evolution equations are known to obey the same equation when effects of dissipation and sources terms are negligible. We investigate the analogy between the two fields for non-vanishing dissipation and sources. In addition to the Reynolds (Re) and Prandtl (Pr_M) numbers, we define a new number (S_M) that is given by the ratio of the diffusive term to the Biermann battery term and which allows for a different classification of magnetised fluid behaviour. Numerical simulations of the two fields are then carried out given a parameter space made of Reynolds, Prandtl and Source numbers. We find it appropriate to present and discuss the findings against Prandtl numbers given that these provide the link between viscous and magnetic diffusion. Our simulations indicate that there exists a range of Prandtl numbers for which the fields remain analogues which raises the important question of how far the analogy goes.

Keywords: *Magnetohydrodynamics, MHD, Magnetic Fields, Fluid Dynamics, Diffusion, Vorticity, Viscosity*

I. INTRODUCTION

Magnetic fields seem to exist everywhere in the universe. The need to explain their presence has led to a number of theories related (i) to their formation (magneto-genesis) or (ii) to their growth (dynamo). One magneto-genesis theory suggests that there were mechanisms that existed that could generate cosmic magnetic fields after, rather than in the Big Bang. Efforts to identify these mechanisms have remained sustained, as is seen in research literature, although an effective mechanism remains elusive. An incomplete list of such efforts may be found in the following articles, review-articles on cosmic magnetic fields [1–3, 5] and in the articles cited in them. Such mechanisms could have operated after recombination during which seed fields of magnitude $B \sim 10^{-20}$ G may have been generated. The Biermann battery [1], which leaves a non-vanishing $\nabla T_e \times \nabla n_e$ (where ∇T_e is the gradient of the temperature of electrons and ∇n_e is the gradient of the electron number density) is an example of a mechanism that can generate magnetic field given the correct conditions.

In the context of the universe, the battery mechanism can operate during the period of galaxy formation and can generate such fields [6, 13]. The fields generated via this mechanism are small in magnitude and therefore there exists a disjuncture between what is observed today which of $B \sim 10^{-6}$ G in magnitude, and what should be observed if no other mechanism operated on the seed field between the time they were generated and today. This is obviously simplistic given our current understanding of the possible evolution of the universe, but therein lies the problem. How did these fields

become so strong? Some amplification mechanisms must act on the seed fields if they are to grow. It seems that different amplification mechanisms, each with a different threshold, are necessary to achieve the present day magnitudes [6]. Magnetic fields have remained difficult to understand firstly because of the technical difficulties encountered in the efforts to observe them, which arises mainly from the fact that these fields are weakness and because of the large distances between the source and the observer. On the other hand the physics of plasma on galactic scales is also not well understood, which accounts for some of the discrepancies between observation and theoretical predictions. These sets of challenges call for a different approach.

We think that *magnetic analogues* are a viable option. Analogues provide an indirect way of studying phenomena that may not readily lend themselves to physical experimentation. A suitable analogue can therefore be a useful tool in providing insight into an otherwise intractable problem. But a complete theory of magnetic analogues is not yet available, although bits and pieces that could make up the theory are strewn in relevant scientific literature. In this article, we examine a correspondence between the magnetic field and fluid dynamics that goes deeper than what is presently available, with the aim of contributing to the searching for a theory of magnetic field analogue. As it has been pointed out elsewhere [17], analogy is not identity. This means that one cannot say that *analogues* are completely equivalent to magnetic fields. Nevertheless such models would be expected to accurately show the key features of magnetic fields. In two articles [19, 20], we examined the analogues of magnetism where the battery terms were necessarily turned off, but not the dissipation. Earlier work on the search for analogues is found in [7–13], where hydrodynamic dynamo models were considered. In this article,

* bob.osano@uct.ac.za

† pw.adams@uct.ac.za

we go beyond hydrodynamics and consider fluids with $\nabla p \neq 0$, with the intention of examining the Biermann battery term effect.

II. RELEVANT ANALOGOUS EQUATIONS

There are two separate systems of equation that concern us: (i) that which includes magnetic field, and (ii) that which includes the vorticity field. We devote this section to the main equations in these systems

II.1. Magnetic induction equation

Our focus is on the modified induction equation, modified in the sense that it carries a source term.

$$\frac{\partial \mathbf{B}}{\partial t} - \nabla \times \mathbf{v} \times \mathbf{B} - \eta \nabla^2 \mathbf{B} = \frac{c \nabla p_e \times \nabla n_e}{n_e^2 e (1 + \chi)}, \quad (1)$$

where $\mathbf{B}$ is the magnetic flux, while η is the coefficient of diffusion. Subscripts e stands for electrons, while p and n are pressure and number density respectively. As pointed out in [6, 22], if one were to assume that the charged fluid is made up three parts: free electrons, protons and hydrogen atoms and where the ionisation fraction (constant in space, for a given temperature) is denoted by χ , it is easy to recast the equation with the setting $p_e = \chi p_{(cf)} / (1 + \chi)$, $n_e = \chi \rho_{(cf)} / m_p$ and $\mathcal{B} = e(1 + \chi)B / m_p$ (other possible parameterisations exist, see for example [21]). The subscript (cf) stands charged fluid. m_p is the proton mass. This conversion yields:

$$\frac{\partial \mathcal{B}}{\partial t} - \nabla \times \mathbf{v} \times \mathcal{B} - \eta \nabla^2 \mathcal{B} = \frac{c \nabla p_{(cf)} \times \nabla \rho_{(cf)}}{\rho_{(cf)}^2}, \quad (2)$$

the form that we will use in the rest of the analysis.

II.2. Fluid vorticity

In this section we briefly review fluid vorticity. Let $\mathbf{u}$ be a vector field describing the motion of neutral fluid. Such a fluid has a vorticity vector $\omega = \nabla \times \mathbf{u}$, which is necessarily solenoidal ($\nabla \cdot \omega = 0$). It is relatively easy to show that the vorticity vector obeys the propagation equation:

$$\frac{\partial \omega}{\partial t} - \nabla \times \mathbf{u} \times \omega - \nu \nabla^2 \omega = - \frac{\nabla p_{(nf)} \times \nabla \rho_{(nf)}}{\rho_{(nf)}^2}. \quad (3)$$

We note that the source terms are comparable if one sets $c = 1$. The subscript nf in Eqn (3) indicates that the fluid is neutral in this case. We use this notation to distinguish it from the charged fluid (cf) given in the previous section. Vorticity is crucial to the understanding of how fluids flow. It will be noted that its importance stems

from the fact that one can recover fluid velocity field by inverting $\omega = \nabla \times \mathbf{u}$ to find an integral over the velocity field, although this is subject to conditions [16]. If the viscosity is negligible, the fluid will have a barotropic equation of state, and the external forces will be conservative and the vorticity will satisfies Helmholtz's laws. These laws necessarily imply that vorticity can not be created and hence the Helmholtz equation:

$$\frac{\partial \omega}{\partial t} = \nabla \times (\mathbf{u} \times \omega) = -\mathbf{u} \cdot \nabla \omega + \omega \cdot \nabla \mathbf{u} - \omega \nabla \cdot \mathbf{u}. \quad (4)$$

It follows that a violation of Helmholtz's laws is necessary for the generation of vorticity. To this end, we consider non-negligible and hence non-barotropic equation of state in our analysis. We also set all external forces to zero for simplicity, where not even the gravitational force is allowed (the alternative will be considered in [24]). We now consider Eqns. (2) and (3) together.

II.3. The analogous system

We investigated and confirmed the well known analogy between magnetic field, $\mathbf{B}$, and vorticity field, ω , equations in [19], wherein we also extended the work to the special case where the magnitude of diffusion equals that of kinetic viscosity. This work has been extended in [20] where the simulation and comparisons have been done for various fluids. It will be noted that these cases studied systems of equations that did not have any source terms. The present study examines the analogy between magnetic field and vorticity field equations in cases where the sources terms are non-negligible and in particular where the Biermann's term is non-vanishing. The relevant system in this case takes the form:

$$\frac{\partial \mathcal{B}}{\partial t} - \nabla \times (\mathbf{v} \times \mathcal{B}) - \eta \nabla^2 \mathcal{B} = \frac{\nabla p_{(cf)} \times \nabla \rho_{(cf)}}{\rho_{(cf)}^2}, \quad (5)$$

$$\frac{\partial \omega}{\partial t} - \nabla \times (\mathbf{u} \times \omega) - \nu \nabla^2 \omega = - \frac{\nabla p_{(nf)} \times \nabla \rho_{(nf)}}{\rho_{(nf)}^2}, \quad (6)$$

where $c = 1$. These equations couple to their corresponding Navier-Stokes equations:

$$\frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} - \nu \nabla^2 \mathbf{v} = - \frac{\nabla p_{(cf)}}{\rho_{(cf)}} \quad (7)$$

$$\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} - \nu \nabla^2 \mathbf{u} = - \frac{\nabla p_{(nf)}}{\rho_{(nf)}} \quad (8)$$

It would appear, from Eqns. (5 and 6) that ω should ideally be compared to $-\mathcal{B}$ which, in practical terms, is matter of the orientation of the field ¹ We proceed

¹ For example, it has been noted in [14] that in some 4 out of 5 galaxies, the radial component of the spiral field could be such that the field points inward. This suggests that the induction equation and the related nonlinearities do not distinguish between $\mathbf{B}$ and $-\mathbf{B}$ other than in the case of Hall effect where direction is important [15]

to simulate and compare various sub-cases of the two systems. Simulations are done using the PENCIL CODE [18] for a periodic box of dimensions 32^3 .

III. SIMULATION RESULTS AND DISCUSSION

There are four different comparisons that we consider. (i) We first look at the magnetic field equations where we compare the evolution of the field with battery term to that without, (ii) this is repeated for the case of vorticity field equations. (iii) We then compare that magnetic and vorticity field equations with battery term, and (iv) without the battery term. These simulations allow us to examine the effect of pressure on the general structure of the magnetic and vorticity fields. It is instructive to note that in the case where pressure is present, pressure terms appear in both the magnetic and vorticity field equations and the respective Navier-Stokes equations to which these field equations are coupled as shown in Eqns.(5-8). We first examine each field individually.

III.1. Evolution of magnetic field

The magnetic induction (hereafter $\mathcal{MI}$) and the Navier - Stokes (hereafter $\mathcal{NS}_{cf}$) equations are coupled, with the level or the strength of coupling heavily depended on the kind of forces at work. For simplicity we choose to neglect the Lorentz and all other external forces, wherein we are interested in the comparative analysis. The Lorentz force is crucial in nonlinear dynamos. It is known that presence of this force causes the dynamo to stop growing because this force changes the flow in such away that the dynamo action is reduced leading to dynamo saturation, but this is not the subject of this article. With vanishing Lorentz force, the coupled system takes the form:

$$\frac{\partial \mathcal{B}}{\partial t} - \nabla \times (\mathbf{v} \times \mathcal{B}) - \eta \nabla^2 \mathcal{B} = \frac{c \nabla p_{(cf)} \times \nabla \rho_{(cf)}}{\rho_{(cf)}^2}, \quad (9)$$

$$\frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} - \nu \nabla^2 \mathbf{v} = -\frac{\nabla p_{(cf)}}{\rho_{(cf)}}. \quad (10)$$

We also use a modified set of this system where pressure has been dropped ($p = 0$) in both equations and wherefore in order to keep track of which system is being simulated, we use the notation $\mathcal{B}_{123}$ to denote Eqn. (9) and $\mathcal{B}_{12}$ to denote the counterpart of this equation that does not have the pressure term (the pressure term is also set to zero in the corresponding $\mathcal{NS}$). We inject an initial seed field of strength 10^{-5} G for the purposes of kickstarting the process. We also set $\eta = \nu = 10^{-10}$ and monitor the *rms* of both $\mathcal{B}_{123}$ and $\mathcal{B}_{12}$. The results of the simulations are given in Fig. (1a).

In order to understand the structure displayed in Fig. (1a), we examine closely the two systems given by Eqns. (

9) and (10) and their pressure-free counterparts. The two growth curves have two points of intersection. The first point of intersection represents the initial equality when only the seed field is present. The seed field then experiences both the amplification due to the dynamo term and the dissipation due to the diffusivity term present. The effect of these two on the magnetic field will differ depending on whether or not the battery term is present. We first consider $\mathcal{B}_{12}$ for a plasma fluid by considering the Reynolds numbers which allow us to compare the relative effects of the two terms on the strength of the magnetic fields. We note that $Re_M = 1$ indicates equality and therefore cancelation of the effects. $Re_M \ll 1$ and $Re_M \gg 1$ signify the perfectly conducting limit (frozen-in flux) and the diffusion equations respectively. For clarity and ease of discussion, let's denote the Reynolds number in this case by $Re_M^{(12)}$ (i.e the Reynolds number for the case where battery term is absent).

So what happens when a source term is present? In this case it is helpful to define a new dimensionless quantity S_M given by the ratio of the source term to the diffusive term (The reader is referred the appendix (A) for a dimensional analysis of this term). The growth pattern of the field can then be assessed based on both the Reynolds and the Source numbers. We now show that in the presence of battery term growth can still be experienced even if $Re_M^{(123)} < 1$ (the Reynolds number for an induction equation that possess a battery term) as long as cumulatively $(Re_M^{(123)} + S_M) > 1$. The evidence for this is found in the simulation results that we will discuss shortly. It is obvious therefore that $(Re_M^{(123)} + S_M) \gg 1$ will lead to a battery-aided dynamo, which is an amendment to the standard dynamo theory. Although these two numbers do allow us to examine the dynamo theory in the context of battery term, the understanding of the actual dynamics requires one to examine the coupling of the induction equation to the Navier-Stokes equation.

In particular, it would appear that the presence of pressure reduces the effect of the dynamo on the field, which can be explained given its effect on the $\mathcal{NS}_{cf}$ Eqn.(10). This reduction is however countered by the contribution from the battery term that increases the strength over and above what was lost via a reduced dynamo effect. The field with the battery term will therefore momentarily have a higher value than the field without. But while the dynamo effect grows because of the change in velocity via the $\mathcal{NS}_{cf}$, the battery term remains constant in time. This means that over time, a point is reached where the effect of the battery term is balanced by the increased dynamo effect and the diffusion i.e $(Re_M^{(12)} = Re_M^{(123)} + S_M)$ which leads to the second point of intersection. From this point on, the reduced effect of dynamo due to the pressure term is much bigger than what can be compensated by the battery term and hence $\mathcal{B}_{12}$ will grow much

faster than $\mathcal{B}_{123}$. It might be interesting to determine how the presence of a battery term modifies the Backus conditions[26] necessary for a dynamo, something that will be pursued in future.

III.2. Evolution of vorticity

As in the case of $\mathcal{MI}$ and $\mathcal{NS}_{cf}$, we consider the evolution of the vorticity equation (hereafter VI) and its corresponding Navier - Stokes equation (hereafter $\mathcal{NS}_{nf}$).

$$\frac{\partial \omega}{\partial t} - \nabla \times (\mathbf{u} \times \omega) - \eta \nabla^2 \omega = - \frac{c \nabla p_{(cf)} \times \nabla \rho_{(cf)}}{\rho_{(cf)}^2}, \quad (11)$$

$$\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} - \nu \nabla^2 \mathbf{u} = - \frac{\nabla p}{\rho}. \quad (12)$$

Here too we use the notation ω_{123} for Eqn. (11) and ω_{12} for the case without the pressure term. The results of the simulations are given in Fig. (1b). The graphs are manifestly different because of the pressure term. We also define two Reynolds numbers ($Re_V^{(12)}$, $Re_V^{(123)}$) and a Source number $\mathcal{S}_V$, the subscript V indicates that these parameters are for the vorticity field system of equations. In this case growth results if $(Re_V^{(123)} - \mathcal{S}_V) \gg 1$

As in the case of magnetic field we find two points of intersections. The first point is the initial equality when only the seed field is present, the second is where the contributions from the various terms balance out leaving the field with a strength similar to that case where the pressure is absent i.e $(Re_V^{(12)} = Re_V^{(123)} - \mathcal{S}_V)$. Although both curves are decaying beyond the second point of intersection, ω_{12} remains stronger than ω_{123} for the same reasons given in the case of magnetic field analysis. In particular, from the $\mathcal{NS}_{nf}$ (12) the pressure leads to a comparatively lower velocity which in turn leads to decreased amplification effect in Eq. (11), this coupled with the reduction on ω_{123} due to the source term ensures that the strength lags that of ω_{12} beyond the second point of intersection. We again warn that the simulation involves weak vorticity and the case of stronger vorticity will be given in a different section.

III.3. Magnetic and Vorticity Induction equations with $Pr_M = 1$ and $Pr_M \neq 1$

We are now at a point where we can compare $\mathcal{MI}$ to $\mathcal{VI}$. Two separate cases; $Pr_M = 1$ ($\nu = \eta = 10^{-5}$) and $Pr_M \neq 1$, will be examined. These systems encode nonlinear dynamics which often times are too complex to study concurrently. For this reason, we will limit the study to special cases and leave extensions and general cases for future studies. It is sufficient to say that nonlinear effects are important in magnetohydrodynamics because they have the capability to induce dynamo action, where magnetic field lines would be stretched by the flow velocity gradients when this exceeds diffusion. Other


nonlinear effects induced by the interaction of a charged with its environment may be analysed using the a novel generalisation of Maxwell's equations and the resulting induction equation presented in [21]. In the present case however, simulation results are given in Figs. (1c) and (1d). Fig.(1c).

The way to understand the curve is to compare them to a straight line with a negative slope, which would be the case were the two to be analogous. We can therefore conclude that given these values for dissipation, $\mathcal{B}_{123}$ approximates a linear function (albeit poorly) better than $\mathcal{B}_{12}$ does. In fact, we see in Fig. (1e) that when we make the dissipation stronger the relationship move progressively towards a straight line and hence better analogue. We have only presented the case for $\mathcal{B}_{123}$, given that the case for $\mathcal{B}_{12}$ will show the same structure although with a much slower progression toward as a straight line. What if these structures are the results of systematic effect in the code and have nothing to with the physical effect with which are associating them with? The way to check this is obtain a plot of the difference $\mathcal{B}_{123}$ and $\mathcal{B}_{12}$ against the difference between ω_{123} and ω_{12} . The idea is that each run will experience the same systematic effects and hence taking the difference eliminates such effects. The results of these simulations are given in Fig.(1d).


If one were to insert the axes, it will be clear that the difference-curves spends its time in the first three quadrants. In the first quadrant, it would appear that the presence of pressure causes difference in the vorticity equations to grow momentarily faster than a comparative growth in the magnetic difference. But the growth in the magnetic difference soon supersedes that of the vorticity and we see a turning, wherein the magnetic difference continues to grow while the vorticity difference declines. The difference in the vorticity return to zero, where the effect of pressure modifies the growth such that the appears to be no difference with the case of pressure-less. The difference then enters the second quadrant where magnetic difference reaches a maximum and begins to decline, while the vorticity difference is now increasingly negative. The magnetic difference then goes to zero where equality is achieved between the pressure-less and pressured equations. From this point on the difference enter the third quadrant where both differences are now both increasingly negative, meaning that the strength of the fields are swapped in both cases. It is conceivable that the graph of such difference for a true analogue would spend its life in the third quadrant. This amounts to eliminating or narrowing the crossover region in Figs. (1a) and (1b). One way of doing this is to scan a parameter space made of $Pr_M \neq 1$. The result of this is given in Fig.(1f), where it is found that analogue status can be achieved for sufficiently small Prandtl numbers. It is known that $Pr_M \ll 1$ means that thermal diffusivity dominates the flow as opposed to $Pr_M \gg 1$ where momentum diffusivity would ordinarily dominate. The importance of the magnetic Prandtl number has also been studied in the context of MHD turbulence in [31]

and [28]


FIG. 1: Simulation results for magnetic and vorticity induction equations involving $Pr_M = 1$ and $Pr_M \neq 1$.




(a) The dotted line represents $\mathcal{B}_{123}$ (with battery term) while the solid line represents $\mathcal{B}_{12}$. (without battery term)




(b) The dotted line represents ω_{123} (with battery term) while the solid line represents ω_{12} . (without battery term)




(c) Again the solid line represents $\mathcal{B}_{123}$ (with battery term) while the dash line represents $\mathcal{B}_{12}$ (without battery term).



(d) The graph represents the plot of the time varying difference $\mathcal{B}_{123} - \mathcal{B}_{12}$ against the time varying difference $\omega_{123} - \omega_{12}$



(e) We have plotted $\mathcal{B}_{123}$ against ω_{123} for different values of ν and η while keeping $Pr_M = 1$. It is apparent that as dissipation becomes stronger the graph tends toward a straight line.



(f) We have plotted $\mathcal{B}_{123}$ against ω_{123} for different values of ν and η while keeping $Pr_M \neq 1$. Here too it is clear that stronger dissipation straightens the curve.

IV. CONCLUSION

In this article we have presented an in depth study of the magnetic and vorticity induction equations. We have defined a new parameter S_M that we call the *Source number* which is given by the ratio of the battery term to the diffusive term, and which will be useful in classifying flows that include source terms. We have analysed the evolutionary behaviours of the two fields with and without the presence of pressure, with the aim of finding constraints that would allow the two to be analogues. We find that, besides the case where the two equations are analogues when dissipation is neglected in the pressure-free case, analogy is still achievable via suitable variation of Prandtl numbers for cases where pressure is non negligible. This in our view strengthens the case for a theory

of magnetic fluid analogues.

V. ACKNOWLEDGEMENT

BO acknowledges URC funding support administered by the University of Cape Town. PA acknowledges funding support from the National Research Foundation (NRF) of South Africa, as well as funding from the University of Cape Town, both administered by the Postgraduate Funding Office (PGFO) of the University of Cape Town.

VI. BIBLIOGRAPHY

-
- [1] Biermann L, *Über den Ursprung der Magnetfelder auf Sternen und im interstellaren Raum* Z. Naturforsch, 5a, 65. (1950)
 - [2] Biermann L and Schlter A, *Cosmic Radiation and Cosmic Magnetic Fields. II. Origin of Cosmic Magnetic Fields*, Phys. Rev. 82, 863 (1951)
 - [3] Durrer R and Neronov A, *Cosmological Magnetic Fields: Their Generation, Evolution and Observation*. The Astronomy and Astrophysics Review, 21, 62. (2013)
 - [4] Subramanian K, *The origin, evolution and signatures of primordial magnetic fields*, arxiv:1504.02311
 - [5] Grasso D and Rubinstein, *Physics Reports, Vol. 348, Issue 3, pp. 163-266*, Physics Reports, Vol. 348, Issue 3, pp. 163-266 (2001)
 - [6] Kulsrud, R M, and Anderson S W, *The spectrum of random magnetic fields in the mean field dynamo theory of the Galactic magnetic field*, ApJ, 396, 606. (1992)
 - [7] Elasser W M, *Induction effects in terrestrial magnetism. Part I. Theory*, Phys. Rev 69, 106 (1946).
 - [8] Elasser W M, *Induction effects in terrestrial magnetism. Part II. Theory*, Phys. Rev 70, 202 (1946).
 - [9] Elasser W M, *Induction effects in terrestrial magnetism. Part III. Theory*, Phys. Rev 72, 821 (1947).
 - [10] Elsasser WM, *The hydromagnetic equations*. Phys Rev 79:183 (1950)
 - [11] Batchelor GK, *On the spontaneous magnetic field in a conducting liquid in turbulent motion*. Proc Roy Soc London A 201:405-416 (1950)
 - [12] Parker E, ApJ 122:293P (1955)
 - [13] Cowling T G, *Magnetic fields of a Sunspots*, M.N., 94, 93.
 - [14] Krause F and Beck R, *Symmetry and direction of seed magnetic fields in galaxies*, Astron. Astrophys. 335 (1998) 789-796.
 - [15] Hollerbach R, Rudiger G, Monthly Notices Roy. Astron. Soc. 337 (2002) 216-224.
 - [16] Saffman P G, *Vortex Dynamics* (Cambridge University Press) 1992
 - [17] Barcelo C, Liberati S and Visser M, *Analogue Gravity*, Living Rev. Relativity, 14, (2011)
 - [18] PENCILCODE, <http://pencil-code.nordita.org>
 - [19] Osano B and Adams P, *Analogue magnetism revisited*, arXiv:1602.09105v1.
 - [20] Adams P and Osano B, *Analogue magnetism: Technical consideration* (2016)
 - [21] Osano B, *Generalised Maxwellian Equations and Implications*, arXiv: 1604.00263.
 - [22] Brandenburg A, Subramanian K, *Astrophysical magnetic fields and nonlinear dynamo theory*, Physics Reports 417 (1), 1-209
 - [23] Malyskin L and Kulsrud R M, *Turbulent Dynamos in Protogalaxies*, Astrophys. J. 571:619-637, 2002
 - [24] Osano B, *Magnetic Analogue in the presence of gravity*, in preparation (2017)
 - [25] Batchelor G K, 1950, *On the spontaneous magnetic field in a conducting liquid in turbulent motion*, Proc. Roy. Soc. London., A201, 405.
 - [26] Backus G, *A class of Self-Sustaining Dissipative Spherical Dynamos*, Annals Phys. 4, 372-447 (1958)
 - [27] Osano, B, *Backus necessary-conditions revisited*, in preparation (2016)
 - [28] Ponty Y and Palitano, *Simulation of Induction at Low Magnetic Prandtl Number*, Phys. Rev. Lett. , 92, 14 (2004)
 - [29] Maxwell J C, *On the physical lines of force*, Philosophical Magazine and Journal of Science, 21 No. 139, pp. 161-175, March 1861.
 - [30] Landau L. D. and Lifshitz E. M. (1987), *Fluid Mechanics*, (2nd ed), Pergamon Press; Landau L.D. and Lifshitz E.M. *The Classical Theory of Fields* (4th ed), Pergamon Press, 1975
 - [31] Shebalin J V, 31st EPS conference on Plasma Phys. London, 28 June-2 July ECA Vol.28G, p-4.203(2004).

VII. APPENDIX

Appendix A: Source Number

In this section, we define a new dimensionless parameter denoted by S_M , which we call the battery number.

It is given by the ratio of the battery term to the diffusion term. The battery term in its original form is $c(k_b \nabla n_e \times \nabla T)/en_e$, where c is the velocity of light (dimension LT^{-1}), k_b the Boltzmann constant (dimension $L^2MT^{-2}\Theta^{-1}$), n_e is the electron number density (dimension; L^{-3}), T is the temperature of the plasma fluid

(dimension Θ), and e is the charge (dimension; IL^{-1}). Writing the variable and constants that make the battery terms in terms of dimensional units, one will find: $ML^2T^{-3}I^{-1}$, which is the same units found when one write the diffusive term in terms of its dimensions. This means that S_M is a dimensionless quantity.